

TILING SPACES ARE INVERSE LIMITS

LORENZO SADUN

ABSTRACT. Let M be an arbitrary Riemannian homogeneous space, and let Ω be a space of tilings of M , with finite local complexity (relative to some symmetry group Γ) and closed in the natural topology. Then Ω is the inverse limit of a sequence of compact finite-dimensional branched manifolds. The branched manifolds are (finite) unions of cells, constructed from the tiles themselves and the group Γ . This result extends previous results of Anderson and Putnam [AP], of Ormes, Radin and Sadun [ORS], of Bellissard, Benedetti and Gambaudo [BBG], and of Gähler [Gäh]. In particular, the construction in this paper is a natural generalization of Gähler's.

1. BACKGROUND

In the last few years, it has become clear that many spaces of tilings of $\mathbb{R}^d$ can be viewed as inverse limit spaces. Anderson and Putnam [AP] began this program for substitution tilings. Given a substitution, they showed that the corresponding space of tilings of $\mathbb{R}^d$ is the inverse limit of a branched d -manifold K under an expansive map from K to itself. If the substitution has a property called “forcing the border” [Kel], then the manifold K is constructed by stitching all the tile types together along possible common boundaries. If the substitution does not force the border, then the construction is similar, only using collared tiles. (A collared tile is a tile that is labeled by the pattern of tiles that touch it). For this construction to work, the tilings must involve only a finite number of tile types (up to translation), meeting full-face to full-face. In particular, the construction does not apply to tilings like the pinwheel [Rad], where tiles appear in an infinite number of orientations.

Ormes, Radin and Sadun [ORS] extended the Anderson-Putnam construction to substitution tilings of $\mathbb{R}^d$ on which the entire Euclidean group acts continuously. Tiles may appear in arbitrary orientations, but there can only be a finite number of tile types *up to Euclidean motion*, and tiles must meet full-face to full-face. The branched manifold has dimension $d(d+1)/2$, which is the dimension of the Euclidean group.

In this construction, a cell in the branched manifold K is not a tile. Rather, a cell is the product of a (possibly collared) tile with $SO(d)$, modulo any (finite!) rotational symmetry that the tile might have. This gives a description of all the ways a tile containing the origin may be placed. The substitution (call it σ) replaces each oriented tile with a union of oriented tiles, giving a map from K to itself. Such a union of tiles is called a *supertile* of order 1. The substitution applied to a supertile of order 1 gives a supertile of order 2, and so on.

1991 *Mathematics Subject Classification*. Primary: 37B50, Secondary: 52C23, 54F65, 37C85.

A point $(x_0, x_1, \dots)$ in the inverse limit $\leftarrow_{\sigma} K$ is a consistent description of a tiling, with x_0 telling how the origin sits inside a tile, x_1 telling how the origin sits inside a supertile of order 1, and x_n telling how the origin sits inside a supertile of order n . If the substitution forces its border (or if we are using collared tiles), the sequence $(x_0, x_1, \dots)$ gives a consistent description of a unique tiling of $\mathbb{R}^d$.

More recently, Gähler [Gäh] and Bellissard, Benedetti and Gambaudo [BBG] have each applied inverse limit methods to tilings that need not be generated by a substitution. If x is a tiling of $\mathbb{R}^d$ that has finitely many tile types *up to translation*, meeting full-face to full-face, then the continuous hull of x (i.e., the closure of the translational orbit of x) is the inverse limit of a sequence of compact branched manifolds $K_0, K_1, K_2, \dots$, under a sequence of maps $\sigma_n : K_n \rightarrow K_{n-1}$, where each branched manifold K_n is the union of (marked) tiles from the original tiling. Of the two constructions, Gähler's is conceptually simpler, but that of Bellissard, Benedetti and Gambaudo appears to be computationally stronger, leading to results such as gap-labeling theorems [BBG].

This paper is an extension of Gähler's construction to tilings of arbitrary Riemannian homogeneous spaces, with general symmetry group. The generalization of the Bellissard-Benedetti-Gambaudo approach to arbitrary spaces is being done independently by Benedetti and Gambaudo [BG].

2. THEOREM AND PROOF

Before stating and proving the result, we must establish some notation. Let M be a Riemannian homogeneous space (such as $\mathbb{Z}^d, \mathbb{R}^d, \mathbf{H}^2, \mathbf{H}^2 \times \mathbb{R}^3$, etc.), and pick a point to be the origin. Let G be the group of isometries of M , let Γ be a closed subgroup of G , and let Γ_0 be the subgroup of Γ that fixes the origin. Let Ω be a collection of tilings of M . We give Ω the topology that two tilings are ϵ -close if they agree on a ball of size $1/\epsilon$ around the origin, up to the action of an ϵ -small element of Γ . We assume that Ω is closed under the action of Γ (i.e., Ω is a union of Γ -orbits), and that Ω is compact. This implies that Ω has finite local complexity, up to the action of Γ .

Theorem. *Ω is the inverse limit of a sequence of compact branched manifolds $K_1, K_2, \dots$ and continuous maps $\sigma_n : K_n \rightarrow K_{n-1}$. The dimension of the branched manifold is the dimension of Γ .*

The idea of the proof is quite simple. A point in the n -th approximant K_n is a description of a tile containing the origin, its nearest neighbors (sometimes called the “first corona”), its second nearest neighbors (the “second corona”) and so on out to the n -th nearest neighbors. (For these purposes, tiles that meet at a point are considered nearest neighbors.) The map $\sigma_n : K_n \rightarrow K_{n-1}$ simply forgets the n -th corona. A point in the inverse limit is then a consistent prescription for constructing a tiling out to infinity. In other words, it is a tiling.

What remains is to actually construct K_n out of geometric pieces and show that K_n is a branched manifold.

First suppose that the tiles are polytopes that meet full-face to full-face. We consider two tiles t_1, t_2 in (possibly different) tilings of M to be equivalent if a patch of the first tiling, containing t_1 and its first n coronas, is identical, up to the action of Γ , to a

similar patch around t_2 . Since Ω has finite local complexity, there are only finitely many equivalence classes, each of which is called an n -collared tile.

For each n -collared tile t_i , we consider how such a tile can be placed around the origin. Let $s_i \subset t_i$ be the set of points where the origin may sit. By finite local complexity, there can only be a finite number of connected components to s_i , and each component is a submanifold of t_i with the same dimension as Γ/Γ_0 . If t_i does not admit any symmetry, then for each point $p \in s_i$, Γ_0 acts simply transitively on the ways to place t_i down with the spot p landing at the origin. The set of ways to place t_i is therefore a principal Γ_0 bundle over s_i , which we denote E_i . The cell $C_i \subset K_n$ associated with t_i is then exactly E_i .

If there are no topological obstructions to trivializing this bundle, we make the identification

$$(1) \quad C_i = E_i = s_i \times \Gamma_0.$$

If M is flat, then there is a *canonical* trivialization of the frame bundle, and this descends to a canonical product (1). If Γ acts transitively on M , then $s_i = t_i$ is contractible, and the decomposition (1), while not canonical, is guaranteed to exist. Although there do exist tilings where neither of these conditions are met, the author knows of no examples where C_i fails to be trivializable.

If t_i admits a discrete symmetry (e.g., is a regular n -gon in a tiling of $\mathbb{R}^2$ or $\mathbf{H}^2$), then more than one point in $s_i \times \Gamma_0$ may describe the same placement of a tile containing the origin. In that case, the cell associated to t_i is the quotient of the Γ_0 bundle E_i by the symmetry. That is,

$$(2) \quad C_i = E_i / \Gamma_{t_i} \quad (= s_i \times_{\Gamma_{t_i}} \Gamma_0, \text{ if } E_i \text{ is trivializable}),$$

where $\Gamma_{t_i} \subset \Gamma_0$ is the group of symmetries of t_i . Since t_i is a collared tile, Γ_{t_i} must be a discrete subgroup of Γ_0 . (Even if a tile had a continuous symmetry, its first corona could not.) By construction, Γ_{t_i} acts without fixed points on E_i , so the interior of C_i is indeed a manifold. (For instance, if $M = \mathbb{R}^2$ and Γ is the Euclidean group, then C_i is a Seifert fibered space. There may be multiple fibers over points of symmetry, but the total space is smooth.)

A patch of a tiling in which the origin is on the boundary of two or more tiles is described by points on the boundary of two or more cells, and these points must be identified. The branched manifold K_n is the disjoint union of the cells C_i , modulo this identification. Since we are using n -collared tiles with $n \geq 1$, each of the points being identified carries complete information about the placement of all the tiles that meet the origin, together with their first $n - 1$ coronas.

We must show that a neighborhood of such a branch point is the union of topological disks whose tangent spaces may be identified. Each such disk is obtained by taking a patch of a tiling in which the above data is actually realized, and considering its orbit under the action of a neighborhood of the identity in Γ . This shows that the dimension of K_n is the dimension of Γ .

Finally, we remove the assumption that the tiles are polytopes that meet full-face to full-face. To a tiling by other shapes we may associate a pattern of marked points, where

a special point is chosen from each tile and labeled by the type of that tile. The Voronoi cells of those points are then polytopes whose faces, properly subdivided, meet full-face to full-face. The original tiling and the tiling by Voronoi cells are mutually locally derivable [BSJ], and so are described by the same topological space, and hence by the same inverse limit structure. ■

In this construction, the group Γ_0 acts naturally on each space K_n , and the maps σ_n are equivariant, from which it follows that

Corollary. *The space Ω/Γ_0 of tilings modulo rotation is the inverse limit of a sequence of compact branched orbifolds K_n/Γ_0 .*

3. EXAMPLES

- (1) If $M = \Gamma = \mathbb{Z}^d$, then we have a $\mathbb{Z}^d$ subshift. The total space Ω is a Cantor set. The n -th approximant K_n is a finite collection of points, corresponding to a decomposition of the Cantor set into a finite number of clopen sets. This decomposition becomes finer as $n \rightarrow \infty$, and the Cantor set is recovered as the inverse limit.
- (2) If $M = \mathbb{R}^d$ and $\Gamma = \mathbb{Z}^d$, then (up to a fixed translation) Ω is a space of tilings of $\mathbb{R}^d$ by square tiles centered at the lattice points. This is a different description of the previous example. In these examples, note that Ω does not have to be the hull of a single tiling, and that the $\mathbb{Z}^d$ action need not be minimal. The $\mathbb{Z}$ subshift on two letters, in which one of the letters appears at most twice, is neither minimal nor the closure of a single orbit, but is an inverse limit space.
- (3) The (d -fold) suspension of a $\mathbb{Z}^d$ subshift has $M = \Gamma = \mathbb{R}^d$. This is a space of tilings of $\mathbb{R}^d$ by unit cubes oriented parallel to the coordinate axes.
- (4) A $\mathbb{Z}^d$ subshift may be suspended in some directions but not in others. For instance, the suspension of a $\mathbb{Z}^2$ subshift in the x direction is a space of tilings of $\mathbb{R}^2$ by square tiles, meeting full-face to full-face, whose centers have integral y coordinate. In this case $M = \mathbb{R}^2$ and $\Gamma = \mathbb{R} \times \mathbb{Z}$.
- (5) The Penrose tiling space, or any other tiling of $\mathbb{R}^d$ with a finite set of prototiles up to translation, has $M = \Gamma = \mathbb{R}^d$. Since Γ_0 is trivial and Γ is the full translation group, the cells C_i can be identified with the collared tiles t_i themselves. This is precisely Gähler's construction. As was shown in [SW], such a space is homeomorphic to the suspension of a $\mathbb{Z}^d$ subshift.
- (6) The pinwheel tiling space [Rad] has $M = \mathbb{R}^2$ and Γ the 2-dimensional Euclidean group [ORS].
- (7) In tiling hyperbolic space, there are a number of interesting choices for Γ . If Γ is a discrete group, then we have the analog of a subshift, associating letters to a discrete set of points in the space being tiled. At the other extreme, one can take Γ to be the entire group of isometries of $\mathbf{H}^n$.
- (8) One dimensional orientable hyperbolic attractors are either solenoids or one dimensional tiling spaces [W, AP]. However, the dyadic solenoid *can* be viewed as a tiling space, of $\mathbf{H}^2$ rather than $\mathbb{R}^1$, following a construction of Penrose [Pen]. See figure 1. In the upper-half-plane model, the basic tile looks like a rectangle, with




FIGURE 1. Penrose's dyadic tiling of hyperbolic space

- the sides of the rectangle geodesics, with the top and bottom edges horocyclic, and with the size chosen such that the bottom edge has twice the length of the top edge. Here the group is $\Gamma = \mathbb{Z} \ltimes \mathbb{R}$, acting on $\mathbf{H}^2$ by $(n, t)(x, y) = (t + 2^n x, 2^n y)$.
- (9) More generally, any geometric substitution in $\mathbb{R}^d$ gives rise to a space of tilings of $\mathbf{H}^{d+1}$, with group $\Gamma = \mathbb{Z} \ltimes \mathbb{R}^d$. As with the dyadic solenoid, it doesn't matter whether the substitution is invertible, since the $\mathbb{Z}$ action enforces the hierarchy. Chaim Goodman-Strauss has adapted this construction to produce a strongly aperiodic set of prototiles for $\mathbf{H}^2$ [G-S1], and to develop a general formalism for describing tilings of hyperbolic space [G-S2].

4. CONCLUSIONS AND OPEN PROBLEMS

The inverse limit structure of Ω implies that the Čech cohomology $H^*(\Omega)$ is the direct limit of $H^*(K_n)$ under the pullback maps σ_n^* . Every element of $H^*(\Omega)$ is the pullback, under the natural projection $\pi_n : \Omega \rightarrow K_n$, of a cohomology class in K_n , for n sufficiently large. If (and only if) $H^*(\Omega)$ is finitely generated, then for n large enough the entire cohomology of Ω is the quotient of $H^*(K_n)$ by the kernel of π_n^* .

To make effective use of this principle, however, requires specific knowledge of the tiling space in question. For substitution tilings, it is easiest to work with the Anderson-Putnam inverse limit construction, rather than that constructed here, although in fact the two are shift equivalent. For cut-and-project tilings with sufficiently nice “windows”, Gähler

[Gäh] has shown that π_n^* is actually an isomorphism in cohomology for n sufficiently large, with the required size of n computable from the geometry of the window.

The inverse limit structure of tiling spaces is related to a possible fiber bundle structure. Locally, Ω looks like a piece of Γ times a Cantor set. Can these neighborhoods be stitched together to yield a fiber bundle (with Cantor set fiber) over a compact manifold? Is that manifold the quotient of the identity component of Γ by a co-compact subgroup? When $M = \Gamma = \mathbb{R}^d$, the answer to both questions is yes [SW], but the general case is not known.

5. ACKNOWLEDGEMENTS

Many thanks to the organizers and participants in the August 2002 conference on Aperiodic Order, Dynamical Systems, Operator Algebras and Topology at the University of Victoria, British Columbia. The results of this paper are a direct consequence of the talks at that conference (especially those of Franz Gähler and Jean-Marc Gambaudo) and the discussions that ensued. Additional thanks to Charles Radin and Michael Baake for assistance with the manuscript. This work was partially supported by the Texas Advanced Research Program.

REFERENCES

- [AP] J. E. Anderson and I. F. Putnam, Topological invariants for substitution tilings and their associated C^* -algebras, *Ergodic Theory & Dynamical Systems* **18** (1998), 509-537.
- [BBG] J. Bellissard, R. Benedetti, and J.-M. Gambaudo, Spaces of tilings, finite telescopic approximations and gap-labelling, Preprint (2001) math.DS/0109062
- [BG] R. Benedetti and G.-M. Gambaudo, On the dynamics of G -solenoids: applications to Delone sets, in preparation.
- [BSJ] M. Baake, M. Schlottmann, and P.D. Jarvis, Quasiperiodic tilings with tenfold symmetry and equivalence with respect to local derivability, *J. Phys. A: Math. Gen.* **24** (1991) 4637-4654.
- [Gäh] F. Gähler, in preparation.
- [G-S1] C. Goodman-Strauss, A strongly aperiodic set of tiles in the hyperbolic plane, preprint, available at <http://comp.uark.edu/~cgstrauss/papers>
- [G-S2] C. Goodman-Strauss, Triangle tilings, preprint 2002.
- [Kel] J. Kellendonk, Noncommutative geometry of tilings and gap labelling, *Rev. Math. Phys.* **7** (1995), 1133-1180.
- [ORS] N. Ormes, C. Radin and L. Sadun, A homeomorphism invariant for substitution tiling spaces, *Geometriae Dedicata* **90** (2002), 153-182.
- [Pen] R. Penrose, Pentaplexity, *Eureka* **39** (1978) 16-22.
- [Rad] C. Radin, The pinwheel tilings of the plane, *Annals of Math.* **139** (1994) 661-702.
- [SW] L. Sadun and R. F. Williams, Tiling spaces are Cantor set fiber bundles, *Ergodic Theory and Dynamical Systems*, in press.
- [W] R.F. Williams, Classification of 1-dimensional attractors, *Proc. Symp. Pure Math.*, **14** (1970), 341-361.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX 78712
E-mail address: sadun@math.utexas.edu